

Rehydration Protocol – Scenario A

- ❑ Calf doesn't drink milk at the AM feed and is scouring, so 2.0L of Osmofit to be offered in AM and lunch or lunch PM (swap for timings)
- ❑ Osmofit (2.0L) can be drenched if the calf has no suck reflex and wont drink (very sick calf), but try in a teated bottle/bucket
- ❑ Calf needs to be offered milk at the PM feed (1-2lt) drenched if no suck reflex

TIME	FEEDING SCHEME	MEAL SIZE
AM h	Osmofit	2.0 L
Lunch time	OsmoFit	2.0 L
PM h	Milk	1-2 L



Rehydration Protocol – Scenario B

- ❑ Calf does drink milk at the AM feed, but doesn't finish and is fussy
- ❑ Osmofit (2.0L) should be offered to the calf in a teated bottle/bucket can be drenched if the calf has no suck reflex and wont drink (very sick calf), but try first in a teated bottle/bucket
- ❑ Calf needs to be offered milk at the PM feed

TIME	FEEDING SCHEME	MEAL SIZE
AM h	Milk	2.0 L
Lunch time	OsmoFit	2.0 L
PM h	Milk	3.0 L



Rehydration Protocol – Scenario C

- ❑ Calf drink milk at the AM feed and then STRESSOR occurs:
High Heat/de horn/Movement/ ANY SICKNESS OR ILL HEALTH
- ❑ Osmofit (2.0L) should be offered to the calf in a teated bottle/bucket between AM/PM milk feeds to combat the STRESSOR
- ❑ Calf needs to be offered milk at the PM feed

TIME	FEEDING SCHEME	MEAL SIZE
AM h	Milk	2.0 L
Lunch time	OsmoFit	2.0 L
PM h	Milk	3.0 L

